

Dienia seidenfadeniana, a new orchid species from Australasia

Dariusz L. SZLACHETKO, Hanna B. MARGOŃSKA & P. RUTKOWSKI

Laboratory of Plant Taxonomy and Phytogeography,
Gdansk University, Al. Legionów 9, PL-80-441 Gdańsk, Poland.
biodarek@univ.gda.pl

KEY WORDS

Dienia,
Orchidaceae,
Australasia.

ABSTRACT

Dienia seidenfadeniana Szlach., Marg. & Rutk., is described and illustrated. The new species appears to be related to *Dienia latifolia* (J.J. Sm.) M.A. Clem. & D.L. Jones, from which it is easily distinguishable by the lip form.

RÉSUMÉ

Dienia seidenfadeniana, une nouvelle espèce d'Orchidaceae d'Australie.

MOTS CLÉS

Dienia,
Orchidaceae,
Australic.

Dienia seidenfadeniana Szlach., Marg. & Rutk. est décrit et illustré. Cette nouvelle espèce est affine de *Dienia latifolia* (J.J. Sm.) M.A. Clem. & D.L. Jones, dont elle se distingue facilement par la forme du labelle.

The genus *Dienia* Lindl. has been recently reinstated by CLEMENTS & JONES (1996). According to these authors it includes 6 species: *Dienia benguetense* (Ames) M.A. Clem. & D.L. Jones, *D. curranii* (Ames) M.A. Clem. & D.L. Jones, *D. latifolia* (J.J. Sm.) M.A. Clem. & D.L. Jones, *D. montana* (J.J. Sm.) M.A. Clem. & D.L. Jones, *D. truncicola* (Schltr.) M.A. Clem. & D.L. Jones and *D. volkensis* (Schltr.) M.A. Clem. & D.L. Jones. It is worthy to note that CLEMENTS & JONES (1996) transferred *Epidendrum ophrydis* J. Koenig to the genus *Crepidium* Blume, despite its similarity to both *Malaxis latifolia* Sm. and *Dienia congesta* Lindl. (GARAY, pers. comm.; SEIDENFADEN, pers. comm.). Both of the latter taxa are treated by SEIDENFADEN (1995) as synonyms of *Malaxis ophrydis* (J. Koenig) Ormerod.

Although the gynostemium structure of *Dienia* Lindl. is similar to *Crepidium* Blume and *Fingardia* Szlach., this genus is characterised by the lip form. Additionally, unlike *Crepidium* and *Fingardia*, the lip of *Dienia* possesses a transverse, rather thick, ridge at the lip base.

While revising the herbarium materials of Malaxidinae from Australasia we found two specimens that in our opinion represent the same unknown taxon that deserves to be separated as a new species. We describe it below.

***Dienia seidenfadeniana* Szlach., Marg. & Rutk., sp. nov.**

Species haec D. latifoliae proxima, sed recedit labello distincte latiori, lobis lateralibus oblique ovato-triangulo-




Fig. 1.—*Dienia seidenfadeniana* Szlach., Marg. & Rutk.: A, part of the pseudobulb; B, flower; C, tepals; D, lip, callus partially removed; E, gynostemium; F, floral bract. (*Bartlett & la Rue 254 (260), L.*).

latibus plus minusve falcatis, lobo mediano fissio et callo trilobularis in parte centrali labelli locato.

TYPUS.—*Bartlett & la Rue 254 (260)*, Sumatra, east coast, Karo-land, Kampong, 2 July 1918 (holo-, L).

PARATYPE.—*van Royen & Sleumer 7508*, Irian Jaya, Vogelkop Peninsula, Alfat River Valley, path from Sururem to Son Village, 720 m, 26 Oct. 1961 (L).

Pseudobulbs 4.5-10 cm long, fusiform, more or less ascending, internodes few to several mm long, 0.2-0.3 cm in diameter. Leaves (2-)9-14, spirally arranged along pseudobulbs; petiole 0.4-2 cm long, narrow, canaliculate, transformed basally into sheath, dark purple; blade 2.2-10.2 cm long, 1.2-2.6 cm wide, oblanceolate to obliquely ovate-lanceolate, acuminate, yellowish-green with purple veins on undersurface. Inflorescence 12.8-17.3 cm long, dark purple, erect, spike 3.8-5.6 cm long, probably longer, 20-60-flowered, dense. Flowers small, non-resupinate, tepals purple with light green edges, lip green. Floral bracts 3.8-6 mm long, linear-lanceolate to linear-triangular, acuminate. Pedicel and ovary 2.5-5 mm long, slender. Dorsal sepal 2-3 mm long, 1-1.6 mm wide, elliptic-ovate to oblong-obovate, obtuse to subobtuse, 3-nerved. Petals 1.8-2.5 mm long, falcately oblong to oblong-oblanceolate, subobtuse to subacute, single- or obscurely 3-nerved. Lateral sepals 2-2.6 mm long, 1.3-1.8 mm wide, obliquely oblong-obovate to oblong-elliptic, subobtuse to subacute, 3-nerved. Lip 2.2-3.4 mm long, 3.2-4 mm wide, distinctly 3-lobed, the basal callus 3-chambered; the middle lobe 0.6-1.4 mm long, 1-1.2 mm wide, ovate to almost triangular in

outline, obtuse, notched in apical third; side lobes falcately ovate-triangular, widely divergent, subacute to subobtuse, with small single tooth near the base of the middle lobe. Gynostemium 1-1.2 mm long, typical for the genus.—Fig. 1.

ETYMOLOGY.—Named in honour of the great Danish orchidologist, Dr. Gunnar SEIDENFADEN.

DISTRIBUTION.—Known so far from Sumatra and Irian Jaya, but probably more widely distributed.

ECOLOGY.—Terrestrial in oak-forest and “deep jungle”.

NOTES.—*Dienia seidenfadeniana* Szlach., Marg. & Rutk. shares the same habit as closely related *D. latifolia*. Both species differ, however, in the lip form. In the newly described entity, the lip is distinctly wider, side lobes of lip obliquely ovate-triangular, more or less falcately divergent, the middle lobe is ovate, split in apical third. The basal lip callus is shallowly 3-chambered.

Acknowledgments

We are grateful to the Curators of BM, HBG, K, P and W for their hospitality during our personal visits, to the Curator of L for the loan of the herbarium materials and to Prof. Dr. Ryszard OCHYRA for the latinization of the diagnoses.

REFERENCES

- CLEMENTS M.A. & JONES D.L. 1996.—*Crepidium myosotis*, a new species of Orchidaceae from Papua New Guinea. *Lasianthera* 1: 32-43.
SEIDENFADEN G. 1995.—*The Descriptiones Epidendrorum of J. G. Koenig, 1791*. Olsen & Olsen, Fredensborg.

*Manuscript received 8 December 1997;
revised version accepted 16 April 1999.*